



Is your electrophoresis good enough for our densitometer?

High Resolution of the SP1809

Scanning gels with the SP1809 is simplicity itself. Permanent and extremely accurate records are made in minutes to be interpreted at leisure. High resolution is




a major feature of this instrument and we supply the test slide to prove it. The SP1809 will separate two lines 0.1 mm thick and 0.1 mm apart (A) and completely resolve two lines 0.2 mm apart (B). Although electrophoretic methods rarely require this degree of resolution, our densitometer is equipped to deal with it.

Convenient Storage and Interpretation

The traces are made and labelled in minutes, but can be interpreted at leisure, easily reproduced and conveniently stored. You can take the eye strain out of gel examination *and* have the research capabilities of the SP1800 to take your analyses much further.

Wide Wavelength Range

Because the SP1809 is incorporated into the advanced SP1800 Spectrophotometer, you get all the advantages of UV-Visible Scanning with a wide wavelength range (nucleic acids and proteins may be detected in the ultraviolet, avoiding lengthy staining and destaining procedures), identification of bands by individual spectra and capability for handling a wide range of sample types.



For full information on the Pye Unicam Electrophoresis System please use the reader reply service or write or phone direct.

Pye Unicam Limited York Street Cambridge England CB1 2PX Telephone Cambridge (0223) 58866 Telex 817331



Pye Unicam Ltd

BIOCHEMICAL SOCIETY SYMPOSIA SERIES

No. 36 NEUROTRANSMITTERS AND METABOLIC REGULATION

Edited by R. M. S. SMELLIE

1972 169 pp. £4.00 (\$12.00)

No. 37 THE STRUCTURE AND FUNCTIONS OF EUKARYOTIC RIBOSOMES

Edited by R. M. S. SMELLIE

1973 130 pp. £3.50 (\$10.50)

No. 38 NITROGEN METABOLISM IN PLANTS

Edited by T. W. GOODWIN and R. M. S. SMELLIE

1973 351 pp. £8.00 (\$20.00)

No. 39 CALCIUM AND CELL REGULATION

Edited by R. M. S. SMELLIE

1974 151 pp. £5.00 (\$13.50)

No. 40 THE METABOLISM AND FUNCTION OF GLYCOPROTEINS

Edited by R. M. S. SMELLIE and J. G. BEELEY

1974 189 pp. £7.00 (\$17.50)

This volume records the proceedings of a symposium held in University College London in April 1974. Each of the ten papers is by an international authority and presents a survey of recent progress in certain aspects of the study of glycoproteins. Topics include methods of analysis, the nature of the carbohydrate-peptide linkage, heterogeneity of glycoproteins, sites of glycosylation, assembly of the carbohydrate moieties, immunoglobulin synthesis, sialic acid modifications, the role of sialic acid and the structure of blood-group-specific glycoproteins in normal and transformed cells.

Available from your bookseller or agent, or direct from

THE BIOCHEMICAL SOCIETY (PUBLICATIONS)
P.O. Box 32, Commerce Way,
Colchester CO2 8HP, Essex

Leaflets giving details of contents of the above publications are available on request

THE BIOCHEMICAL SOCIETY
SPECIAL PUBLICATIONS SERIES

No. 1 BIOCHEMISTRY AND MENTAL ILLNESS

Edited by L. L. IVERSON and S. P. R. ROSE

1973 251 pp. £4.50 (\$12.00)

No. 2 BIOCHEMISTS IN INDUSTRY

Edited by K. W. FULLER

1973 58 pp. £1.25 (U.S. \$3.50)

No. 3 THE METABOLISM AND FUNCTION OF VITAMIN D

Edited by D. R. FRASER

1974 114 pp. £4.00 (\$10.50)

**No. 4 MEMBRANE ADENOSINE TRIPHOSPHATASES AND
TRANSPORT PROCESSES**

Edited by J. R. BRONK

1974 197 pp. £5.00 (\$12.50)

Available from your bookseller or agent, or direct from

THE BIOCHEMICAL SOCIETY (PUBLICATIONS)
P.O. Box 32, Commerce Way,
Colchester CO2 8HP, Essex

Leaflets giving details of contents of the above publications are available on request

ADVERTISING RATES

| | |
|--------------------------------|-----|
| Cover and Special Positions | £90 |
| Full Page | £70 |
| Half Page | £40 |
| Quarter Page | £25 |
| Agency Commission | 10% |
| Publisher's Discount | 10% |

All communications regarding advertising matters should be addressed to:

Advertisement Section,
Biochemical Society,
7 Warwick Court,
London WC1R 5DP
Telephone: 01-242 1076
(4 lines)

THE BIOCHEMICAL JOURNAL

The Biochemical Journal Cumulative Indices

Save Research Time

Busy researchers and authors find that these multi-volume indices are invaluable. Where time is money the cost is recovered quickly. Produced as a service to subscribers and priced accordingly.

| Volumes | £ p | U.S.\$ |
|---------|------|--------|
| 21-30 | 0.75 | 2.50 |
| 31-40 | 2.00 | 5.50 |
| 41-65 | 2.75 | 7.50 |
| 66-90 | 3.25 | 9.00 |
| 91-115 | 5.25 | 13.00 |
| 116-130 | 5.50 | 15.00 |

All prices include postage (by surface mail to overseas)

Available from:

The Biochemical Society
(Publications)

PO BOX 32 Commerce Way
Whitehall Industrial Estate
Colchester CO2 8HP Essex

CLINICAL SCIENCE AND MOLECULAR MEDICINE

Formerly *Clinical Science*

EDITORIAL BOARD

For the Biochemical Society

J. A. OWEN, *Deputy Chairman*

B. H. BILLING, A. M. BRECKENRIDGE, C. N. HALES, T. D. R. HOCKADAY,
R. G. HUNTSMAN, G. H. LATHE, J. LIDDELL, I. MACINTYRE,
K. L. MANCHESTER, R. W. E. WATTS

For the Medical Research Society

R. D. COHEN, *Chairman*

G. CUMMING, D. C. FLENLEY, N. F. JONES, D. M. MATTHEWS,
M. E. M. NOBLE, D. K. PETERS, A. POLAK, J. I. S. ROBERTSON,
B. ROBINSON, J. D. M. SLATER

VOL. 48, No. 4

APRIL 1975

CONTENTS

- Correlation of peak aortic and carotid flow acceleration during coronary occlusion in conscious baboons. By J. E. CHIMOSKEY, W. J. FLANAGAN, L. L. HUNTSMAN, E. GAMS and R. F. RUSHMER
- Triglyceride, free fatty acid and carbohydrate metabolism in hyperlipaemic (type IV) and normolipaemic subjects on carbohydrate- or fat-rich diets. By L. HAGENFELDT, K. HELLSTRÖM and J. WAHREN
- Enzyme activities and properties of lysosomes and brush borders in jejunal biopsies from control subjects and patients with coeliac disease. By T. J. PETERS, J. R. HEATH, M. H. WANSBROUGH-JONES and W. F. DOE
- Assessment of urine-concentrating ability in man: effect of fludrocortisone and urea in enhancing response to vasopressin. By C. de F. W. GOONARATNA and O. M. WRONG
- Ammonia and urea transport by the excluded human colon. By R. L. BOWN, J. A. GIBSON, J. C. B. FENTON, W. SNEDDEN, M. L. CLARK and G. E. SLADEN
- Human platelet monoamine oxidase activity in iron-deficiency anaemia. By MOUSSA B. H. YOUNDIM, H. F. WOODS, B. MITCHELL, D. G. GRAHAME-SMITH and SHEILA CALLENDER
- Variation in the proportions of individual pepsins secreted by the cat in response to vagal stimulation and hypoglycaemia. By CHRISTINE L. WRIGHT, B. SHAW, D. J. SANDERS and J. D. REED
- Effect of bile-duct ligation on organelle marker enzymes in the liver and serum of rats. By T. J. PETERS, G. NEALE and J. R. HEATH
- The selective inhibition of chenodeoxycholate synthesis by cholate metabolites in man. By E. W. POMARE and T. S. LOW-BEER
- SHORT COMMUNICATIONS
- The effect of age on deoxyribonucleic acid synthesis in rat adipose tissue. By B. M. LEWIS and T. M. HAYES
- Role of saline consumption in the prevention of deoxycorticosterone hypertension in rats by central 6-hydroxydopamine. By P. J. LEWIS, H. J. DARGIE and C. T. DOLLERY
- Postural fluid retention in patients with idiopathic oedema: lack of relationship to the phase of the menstrual cycle. By O. M. EDWARDS and R. I. S. BAYLISS

*Subscription: £2.00 (\$7.50) per part; £20.00 (\$70.00) per year.
Orders may be placed with your bookseller, or sent direct to the publishers.*


BLACKWELL SCIENTIFIC PUBLICATIONS LTD

5 ALFRED STREET, OXFORD OX1 4HB, ENGLAND



Tetramethylbenzidine

A safe, specific and sensitive reagent for the detection of blood



For many years benzidine has been used as a sensitive and specific reagent for the detection of blood.¹ However, its extreme carcinogenicity has curtailed its use in recent years and prompted the search for a safe substitute. It has been shown that the *o*-hydroxy substituent contributes to the carcinogenic activity of aromatic amines. Thus, 3,3'-dihydroxybenzidine, a metabolite of benzidine, has been found to be more carcinogenic than benzidine itself. Since 3,3'-dimethylbenzidine was found to be a sensitive reagent for the detection of blood and less harmful than benzidine, it seemed likely that 3,3',5,5'-tetramethylbenzidine, in which *o*-hydroxylation is impossible, would be a sensitive and safe substitute for benzidine. Indeed, subcutaneous injection of 3,3',5,5'-

tetramethylbenzidine into rats "produced no tumors specifically attributable to it, in doses greater than those in which benzidine or *o*-tolidine cause a high yield of neoplasms."²

Blood has been found to exhibit a peroxidase-like activity, yielding, in the presence of hydrogen peroxide, colored products with certain substrates, e.g., benzidine. 3,3',5,5'-Tetramethylbenzidine proved to be as sensitive as benzidine in routine clinical tests for occult blood in urine and feces.² The sensitivity, specificity and ease-of-handling make it a useful reagent for the detection of blood for forensic purposes. 3,3',5,5'-Tetramethylbenzidine is potentially useful in biological redox systems.

References:

1. E.C. Miller, J.A. Miller and H.A. Hartman, *Cancer Res.*, **21**, 815 (1961).
2. V.R. Holland, B.C. Saunders, F.L. Rose and A.L. Walpole, *Tetrahedron*, **30**, 3299 (1974).

86,033-6

3,3',5,5'-Tetramethylbenzidine

100mg \$18.00

Aldrich Chemical Company, Inc.

Craftsmen in Chemistry

Corporate Offices:
Aldrich Chemical Co., Inc.
940 W. Saint Paul Ave.
Milwaukee, Wisconsin 53233

In Great Britain:
Aldrich Chemical Co., Ltd.
264 Water Rd., Wembley, Middx.
HAO 1PY, England

In Continental Europe:
Aldrich-Europe
B-2340 Beerse
Belgium

In Germany:
EGA-Chemie KG
7924 Steinheim am Albuch
Germany