



More "FIRSTS" from Sigma

DIHYDROFOLATE REDUCTASE

(Tetrahydrofolate Dehydrogenase, 5, 6, 7, 8-Tetrahydrofolate: NADP oxidoreductase; E.C. No. 1.5.1.3)

D 6385 – Type II From Bovine Liver Ammonium Sulfate Suspension

0.5 unit £ 5.18 5 units £ 28.75
1 unit 8.63 25 units . 95.84

ACTIVITY: Approx. 8 U/mg Protein (W/C)

Suitable for assay of Dihydrofolic Acid per Matthews and Heunneken, JBC 238:3436, (1963)

UNIT DEFINITION: One unit will convert 1.0 μ Mole of 7, 8-Dihydrofolate and NADPH to 5, 6, 7, 8-Tetrahydrofolate and NADP per minute at pH 6.5 at 25°C.

5, 10-METHYLENE TETRAHYDROFOLATE DEHYDROGENASE

(5, 10-Methylenetetrahydrofolate: NADP oxidoreductase; E.C. No. 1.5.1.5)

M 4882 – Type III From Yeast Ammonium Sulfate Suspension

0.5 unit £ 7.28 5 units £ 40.31
1 unit 12.09 10 units 67.19

ACTIVITY: 0.5 to 1.0 U/mg Protein (W/C)

Suitable for the assay of Tetrahydrofolic Acid per Ramasastri and Blakley, JBC 239:106, (1964)

UNIT DEFINITION: One unit will convert 1.0 μ Mole of 5, 10-Methylenetetrahydrofolic Acid and NADP to 5, 10-Methenyltetrahydrofolic Acid and NADPH per minute at pH 7.5 at 25°C.

We also offer...

FORMIMINO-L-GLUTAMIC ACID (FIGLU) ENZYMES

No. 365-2 From Porcine Liver

A crude mixture of Formimino-L-Glutamic Acid Transferase and Formimino-THF-cyclodeaminase.

1.5 units £ 4.84
3.0 units 8.06
6.0 units 13.44

ACTIVITY: 0.2 to 0.5 U/mg.

Suitable for the Determination of Formimino-L-Glutamic Acid (FIGLU) in Urine or other Fluids, per Sigma Technical Bulletin No. 365.

UNIT DEFINITION: One unit will convert 1.0 μ Mole of FIGLU and THF to 5, 10-Methenyl-THF per minute at pH 7.2 at 25°C.

Ref: Tabor and Wyngarden, *J. Clin. Invest.* 37, 824, (1958)

FORMIMINO-L-GLUTAMIC ACID (FIGLU)

No. F 8626 – Hemi-Barium Salt

10 mg £ 0.96 100 mg £ 5.70
25 mg 1.87 250 mg 11.98
50 mg 3.35 1 g 45.52

No. 365-10 – Standard solution 2 ml £ 1.20

Contains 2.0 μ Moles FIGLU per ml.
For the Determination of FIGLU in Urine or other applications.

dl-N-5-METHYL TETRAHYDROFOLIC ACID

DIHYDROFOLIC ACID FOLIC ACID

9/75

dl-L-TETRAHYDROFOLIC ACID

ORDER DIRECT

TELEPHONE COLLECT



from ANYWHERE in Great Britain

or Europe, 01-549-3171

— ● —
Night and Holidays

N. A. Virgo 01-942-0883

— ● —
Most orders will be shipped within hours

CABLE: SIGMALOND, London, England

It's a Pleasure Doing Business With Sigma!

SIGMA LONDON

CHEMICAL COMPANY LTD.

Norbinton Station Yard, Kingston-upon-Thames, Surrey, KT2 7BH, England

MANUFACTURERS OF THE FINEST BIOCHEMICALS AVAILABLE

Home office:

SIGMA Chemical Co.—P.O. Box 14508—St. Louis, Missouri, 63178 U.S.A.
Telephone: 314-771-5750 (Reverse Charges)

Also distributed through:

SIGMA Chemie GmbH MÜNCHEN • D-8014-Neubiberg, Isarstrasse 14, W. Germany
Telephone: 089/60 70 03 (Reverse Charges)

NEW FROM MILES LABORATORIES, INC.

CHOLESTEROL ESTER HYDROLASE

The enzymatic determination of serum cholesterol has recently been reported (1,2). This method involves the hydrolysis of cholesterol esters to free cholesterol and fatty acids. Cholesterol is then oxidized to cholest-4-en-3-one with the production of hydrogen peroxide. Hydrogen peroxide production is determined enzymatically and followed spectrophotometrically. Cholesterol ester hydrolase and cholesterol oxidase are crucial elements of this system.

Miles Research Products is pleased to offer cholesterol ester hydrolase extracted from bovine pancreas by a method developed in our laboratories. Our preparation has a CEH activity of approximately 0.2 units per milligram of material. It is supplied as a freeze-dried powder containing about 45% phosphate buffer salts and about 20% sodium cholate. The residual proteolytic enzyme content has been specifically reduced to yield a stable product. Further information and assay details are available on request.

Code No.	Pkg. size	Price
36-129-2	100 un	\$ 30.00
36-129-3	1000 un	200.00
36-129-4	5000 un	800.00

1) Allain, et al., (1974), *Clinical Chemistry*, 20, 470.

2) Tarbutton and Gunter, (1974), *Clinical Chemistry*, 20, 724.

RESEARCH PRODUCTS



Research Products
Miles Laboratories, Inc.
Elkhart, Indiana 46514
Phone: 219-264-8804

Miles Laboratories, Ltd.
Post Office Box 37, Stoke Poges
Slough, England SL 2 4 LY
Phone: Farnham Common 2151

The
Centre
for
radiochemicals

Cyclic nucleotides:

improve the sensitivity of your assays with these super-hot compounds

[2,8-³H] adenosine 3',5'-cyclic phosphate TRK.498
current batch at 36 Ci/mmol

[8-³H] guanosine 3',5'-cyclic phosphate TRK.392
current batch at 19 Ci/mmol



Full details on request.

**The Radiochemical Centre
Amersham**

The Radiochemical Centre Limited, Amersham, England.
Tel: Little Chalfont (024 04) 4444
In the Americas: Amersham/Searle Corp, Illinois 60005. Tel: 312-593-6300.
In W.Germany: Amersham Buchler GmbH & Co, KG, Braunschweig.
Tel: 05307-4693-97

0442

Hoppe-Seyler's Zeitschrift für Physiologische Chemie

Begründet 1877 · Fortgeführt von A. Kossel, F. Knoop und K. Thomas

Editors in Chief:

A. BUTENANT · F. LYNEN · G. WEITZEL

Subscription Rates;

For one volume (12 parts) DM 480,—

Vol. 356 No. 8

Contents

August 1975

On the specificity and stereospecificity of the conversion of different 2,3-unsaturated acid by *Clostridium kluveri*

H. Hashimoto, H. Günther and H. Simon

The occurrence of a reductase of Δ^2 -Carboxylic acids in *Clostridium kluveri* with a stereospecificity different from that of butyryl-CoA-dehydrogenase

H. Hashimoto, B. Rambeck, H. Günther, A. A. Mannschreck and H. Simon

Free flow electrophoresis, I: Theoretical and experimental investigations of the influence of mechanical and electrokinetic variables on the efficiency of the method

K. Hannig, H. Wirth, B.-H. Meyer and K. Zeiller

Free-flow electrophoresis, II: Analysis of the method with respect to preparative cell separation

K. Zeiller, R. Löser, G. Pascher and K. Hannig
Subcellular distribution and properties of kynurenine pyruvate transaminase in rat kidney

T. Noguchi, M. Nakatani, Y. Minatogawa, M. Morimoto and R. Kido

Purification and properties of cytochrome *c*-556 from *Agrobacterium tumefaciens* B2a

C. Van den Branden, J. Van Beeumen, J. De Ley and A. Van de Mierop

Formation of a complex between yeast proteinases A and B

H. Hünze, H. Betz, T. Saheki and H. Holzer

Glucosidases specific for the cyanogenic glucoside triglochinin. Purification and characterization of β -glucosidases from *Alocasia macrorrhiza* Schott

W. Hösel and A. Nahrstedt

Activity and isoenzymes of lactate dehydrogenase in tissues of amphibians (*Xenopus laevis*, *Ambystoma mexicanum*, *Triturus alpestris* and *vilgaris*) and the response in the fat body of *Xenopus* males to gonadotropin injection

I. Faulhaber and L. Lyra

Microelectrophoresis in continuous polyacrylamide gel gradients, IV. The effect of reducing agents on the electrophoresis of proteins in sodium dodecylsulfate buffer systems

R. Rüchel, C. Richter-Landsberg and V. Neuhoff

Extraction of collagenase from the 6000 \times g sediment of uterine and skin tissues of mice. A comparative study

G. Wirl

Synthesis of a chemically reactive analog of the nonsense codon U-G-A

Its reaction with ribosomes of *Escherichia coli*

O. Pongs and E. Rössner

Studies on polypeptides, VI: Synthesis, circular dichroism and immunological studies of tyrosyl C-peptide of human proinsulin

V. K. Naithani, M. Dechesne, J. Markussen, L. G. Heding and U. D. Larsen

Relevance of three-dimensional reconstructions of stain distributions for structural analysis of biomolecules

W. Baumeister and M. Hahn

Comments on the paper "Relevance of three-dimensional reconstructions of stain distributions for structural analysis of biomolecules"

W. Hoppe, J. Gaßmann, N. Hunsmann, H. J. Schramm and M. Sturm

Short Communications

Countercurrent distribution of polysulfonated proteins

G. Braunitzer, S. J. van der Walt and B. Bless

Investigation of the proposed anhydride intermediate in the asparaginase reaction

V. M. Bieluch and K. Hellman

Influence of membrane-bound *N*-acetyl-neuraminic acid on the survival of erythrocytes in man

J. Jancik, R. Schauer and H.-J. Streicher

The primary structure of a monoclonal IgA-Immunoglobulin (IgA Tro.), I: The amino acid sequence of the L-chain of λ -type, subgroup II

R. Scholz and N. Hilschmann

The primary structure of a monoclonal IgA-Immunoglobulin (IgA Tro.), II: The amino acid sequence of the H-chain, α -type, subgroup III; structure of the complete IgA-molecule

H. Kratzin, P. Altevogt, E. Ruban, A. Kortt, K. Staroscik and N. Hilschmann

Indexed in Current Contents



Walter de Gruyter · Berlin · New York

ADVERTISING RATES

Cover and Special Positions	£90
Full Page	£70
Half Page	£40
Quarter Page	£25
Agency Commission	10%
Publisher's Discount	10%

All communications regarding advertising matters should be addressed to:

Advertisement Section,
Biochemical Society,
7 Warwick Court,
London WC1R 5DP
Telephone: 01-242 1076
(4 lines)

THE BIOCHEMICAL JOURNAL

The Biochemical Journal Cumulative Indices

Save Research Time

Busy researchers and authors find that these multi-volume indices are invaluable. Where time is money the cost is recovered quickly. Produced as a service to subscribers and priced accordingly.

Volumes	£ p	U.S.\$
21-30	0.75	2.50
31-40	2.00	5.50
41-65	2.75	7.50
66-90	3.25	9.00
91-115	5.25	13.00
116-130	5.50	15.00

All prices include postage (by surface mail to overseas)

Available from:

The Biochemical Society
(Publications)

PO BOX 32 Commerce Way
Whitehall Industrial Estate
Colchester CO2 8HP Essex

CLINICAL SCIENCE AND MOLECULAR MEDICINE

Formerly *Clinical Science*

EDITORIAL BOARD

For the Biochemical Society

J. A. OWEN, *Deputy Chairman*

B. H. BILLING, A. M. BRECKENRIDGE, C. N. HALES, T. D. R. HOCKADAY,
R. G. HUNTSMAN, G. H. LATHE, J. LIDDELL, I. MACINTYRE,
K. L. MANCHESTER, R. W. E. WATTS

For the Medical Research Society

R. D. COHEN, *Chairman*

G. CUMMING, D. C. FLENLEY, N. F. JONES, D. M. MATTHEWS,
M. E. M. NOBLE, D. K. PETERS, A. POLAK, J. I. S. ROBERTSON,
B. ROBINSON, J. D. M. SLATER

VOL. 49, No. 4

OCTOBER 1975

CONTENTS

- Paradoxical diuresis after vasopressin administration to patients with neurohypophyseal diabetes insipidus treated with chlorpropamide, carbamazepine or clofibrate. By A. E. MEINDERS, A. M. VAN LEEUWEN, J. G. G. BORST and V. CEJKA
- Plasma catecholamines, plasma renin activity and plasma aldosterone in tetraplegic man horizontal and tilted. By C. J. MATHIAS, N. J. CHRISTENSEN, J. L. CORBETT, H. L. FRANKEL, T. J. GOODWIN and W. S. PEART
- Increased plasma calcitonin in early acute renal failure. By R. ARDAILLOU, M. BEAUFILS, MARIE-PAUL NIVEZ, R. ISAAC, C. MAYUAD and J.-D. SRAER
- Evidence for active transport of tripeptides by hamster jejunum *in vitro*. By JILL M. ADDISON, D. BURSTON, J. W. PAYNE, S. WILKINSON and D. M. MATTHEWS
- A common mechanism for transport of di- and tri-peptides by hamster jejunum *in vitro*. By JILL M. ADDISON, D. BURSTON, JUDITH A. DALRYMPLE, D. M. MATTHEWS, J. W. PAYNE, M. H. SLEISENGER and S. WILKINSON
- Body weight and body water in chronic cor pulmonale. By R. H. A. CAMPBELL, HILARY L. BRAND, J. R. COX and P. HOWARD
- Cardiac effects of lead in drinking water of rats. By M. R. MOORE, P. A. MEREDITH, A. GOLDBERG, K. E. CARR, P. G. TONER and T. D. V. LAWRIE
- Effect of phenylalanine and its metabolites on the metabolism of leucocytes and lymphocytes. By D. M. BRADLEY and R. F. MAHLER
- Effect of intravenous frusemide on plasma renin concentration: suppression of response in hypertension. By P. L. PADFIELD, M. E. M. ALLISON, J. J. BROWN, A. F. LEVER, R. G. LUKE, C. C. ROBERTSON, J. I. S. ROBERTSON and M. TREE
- Defective membrane systems in dystrophic skeletal muscle of the UM-X7.1 strain of genetically myopathic hamster. By N. S. DHALLA, A. SINGH, S. L. LEE, M. B. ANAND, A. M. BERNATSKY and G. JASMIN

Subscription: £2.00 (\$7.50) per part; £20.00 (\$70.00) per year.

Orders may be placed with your bookseller, or sent direct to the publishers.

BLACKWELL SCIENTIFIC PUBLICATIONS LTD

5 ALFRED STREET, OXFORD OX1 4HB, ENGLAND

NO HIDDEN EXTRAS
MANY EXTRAS INCLUDED

**LATEST
ADDITIONS**

Boehringer Mannheim

Benefits

for you

- Confidence through Quality Control — probably the most exacting in the world
- Service — first class, unequalled
- Economy — cost competitive especially when related to performance
- Innovation — constant developments through progressive research

A few of our latest examples

Preparation	Pack Size	Cat. No.
Acetyl-CoA synthetase	5 mg (20 U)	15466
Adenosine-5'-O-thiomonophosphate	25 mg	15102
Adenylyl-(3'-5')-cytidine	10 mg	15145
Carboxypeptidase C	1 U (37°C)	15662
Clostripain	20 U	15061
Cyclic G.M.P. Test Combination with G3.5MP binding protein	100 tests	15266
Di (adenosine-5'-)pentaphosphate (Ap5A)	25 mg	15398
Succinyl-CoA synthetase	5 mg (50 U)	15062
Valinomycin	25 mg	15556

For full details
of biochemicals
please contact:



The Boehringer Corporation (London) Ltd.,
Bilton House, Uxbridge Road, London W5 2TZ.
Telephone: 01-679 6944 (7 lines) Telex: 934289
or in countries other than Gr. Britain your
local Boehringer Mannheim Office.