

**Reviews**

The toxicity *in vitro* of  $\beta$ -amyloid protein  
by L.L. Iversen, R.J. Mortishire-Smith, S.J. Pollack and M.S. Shearman 1-16

Serine/threonine protein phosphatases  
by S. Wera and B.A. Hemmings 17-29

**Research Communications**

Phosphatidylinositol 3-kinase activity is required for early endosome fusion A.T. Jones and M.J. Clague 31-34

Physiological control of metabolic flux: the requirement for multisite modulation D.A. Fell and S. Thomas 35-39

Putative capacitative calcium entry channels: expression of *Drosophila trp* and evidence for the existence of vertebrate homologues C.C.H. Petersen, M.J. Berridge, M.F. Borgese and D.L. Bennett 41-44

The secreted form of the Alzheimer's  $\beta$ -amyloid precursor protein stimulates a membrane-associated guanylate cyclase S.W. Barger and M.P. Mattson 45-47

**Research Papers**

**Proteins**

A novel lipid-binding protein from the cestode *Moniezia expansa* D. Janssen and J. Barrett 49-57

Mass-spectrometric analysis of ADP-ribosylation factors from bovine brain: identification and evidence for homogeneous acylation with the C<sub>14:0</sub> fatty acid (myristate) S.J. Berger, K.A. Resing, T.C. Taylor and P. Melançon 125-132

The refolding of hen egg white riboflavin-binding protein: effect of protein disulphide isomerase on the reoxidation of the reduced protein D.A. McClelland, S.H. McLaughlin, R.B. Freedman and N.C. Price 133-137

Magnetic circular dichroism study of the selenium-substituted form (Fe<sub>3</sub>Se<sub>4</sub>) of bovine heart aconitase J.L. Breton, J.A. Farrar, M.C. Kennedy, H. Beinert and A.J. Thomson 197-202

Demonstration by pulsed neutron scattering that the arrangement of the Fab and Fc fragments in the overall structures of bovine IgG1 and IgG2 in solution is similar M.O. Mayans, W.J. Coadwell, D. Beale, D.B.A. Symons and S.J. Perkins 283-291

Molecular cloning of mouse intestinal trefoil factor and its expression during goblet cell changes M. Tomita, H. Itoh, N. Ishikawa, A. Higa, H. Ide, Y. Murakumo, H. Maruyama, Y. Koga and Y. Nawa 293-297

Lysosomal processing of amyloid precursor protein to A $\beta$  peptides: a distinct role for cathepsin S J.S. Munger, C. Haass, C.A. Lemere, G.-P. Shi, W.S.F. Wong, D.B. Teplow, D.J. Selkoe and H.A. Chapman 299-305

Identification of histatins as tannin-binding proteins in human saliva Q. Yan and A. Bennick 341-347

**Enzymes**

Cellobiohydrolase B, a second exo-cellobiohydrolase from the cellulolytic bacterium *Cellulomonas fimi* H. Shen, N.R. Gilkes, D.G. Kilburn, R.C. Miller, Jr. and R.A.J. Warren 67-74

Heterologous processing of rat prosomatostatin to somatostatin-14 by PC2: requirement for secretory cell but not the secretion granule A.S. Galanopoulou, N.G. Seidah and Y.C. Patel 111-118

A novel phospholipase A<sub>2</sub> from human placenta W.-J. Buhl, L.M. Eisenlohr, I. Preuss and U. Gehring 147-153

Regulation of the  $\beta$ -lactamase BlaL of *Streptomyces cacaoi*: the product of the *blaB* regulatory gene is an internal membrane-bound protein J. Magdalena, B. Joris, J. Van Beeumen, R. Brasseur and J. Dusart 155-160

Evidence for an essential serine residue in the active site of the Theta class glutathione transferases P.G. Board, M. Coggan, M.C.J. Wilce and M.W. Parker 247-250

Characterization by spectroscopic, kinetic and equilibrium methods of the interaction between recombinant human cystatin A (stefin A) and cysteine proteinases E. Pol, S.-L. Olsson, S. Estrada, T.W. Prasthofer and I. Björk 275-282

Expression, purification and characterization of recombinant human *N*-acetylgalactosamine-6-sulphatase J. Bielicki, M. Fuller, X.-H. Guo, C.P. Morris, J.J. Hopwood and D.S. Anson 333-339

**Carbohydrates and lipids**

Identification and subcellular localization of sphinganine-phosphatases in rat liver P. De Ceuster, G.P. Mannaerts and P.P. Van Veldhoven 139-146

Heparin proteoglycans synthesized by mouse mastocytoma contain chondroitin sulphate	K. Lidholt, I. Eriksson and L. Kjellén	<b>233–238</b>
The biosynthesis of GDP-D-arabinopyranose in <i>Crithidia fasciculata</i> : characterization of a D-arabino-1-kinase activity and its use in the synthesis of GDP-[5- <sup>3</sup> H]D-arabinopyranose	P. Schneider, A. Nikolaev and M.A.J. Ferguson	<b>307–315</b>
<b>Gene structure and expression</b>		
Expression of stress proteins and mitochondrial chaperonins in chronically stimulated skeletal muscle	O.I. Ornatsky, M.K. Connor and D.A. Hood	<b>119–123</b>
Reduced expression of <i>kan-1</i> (encoding putative bile acid-CoA-amino acid <i>N</i> -acyltransferase) mRNA in livers of rats after partial hepatectomy and during sepsis	M. Furutani, S. Arai, H. Higashitsuji, M. Mise, M. Fukumoto, S. Takano, H. Nakayama, M. Imamura and J. Fujita	<b>203–208</b>
Characterization of bovine tracheobronchial phenol sulphotransferase cDNA and detection of mRNA regulation by cortisol	S.J. Schauss, T. Henry, R. Palmatier, L. Halvorson, R. Dannenbring and J.D. Beckmann	<b>209–217</b>
An unusual arrangement of 13 zinc fingers in the vertebrate gene <i>Z13</i>	T.C. Schulz, B. Hopwood, P.D. Rathjen and J.R.E. Wells	<b>219–224</b>
Identification of a functional promoter element in the 5'-flanking region of the rat cMG1/TIS11b gene	A.N. Corps, J.C. Pascall, K.M. Hadfield and K.D. Brown	<b>251–258</b>
<b>Regulation of metabolism</b>		
Failure of insulin-regulated recruitment of the glucose transporter GLUT4 in cardiac muscle of obese Zucker rats is associated with alterations of small-molecular-mass GTP-binding proteins	I. Uphues, T. Kolter, B. Goud and J. Eckel	<b>161–166</b>
Modulation of hepatic apolipoprotein B, 3-hydroxy-3-methylglutaryl-CoA reductase and low-density lipoprotein receptor mRNA and plasma lipoprotein concentrations by defined dietary fats. Comparison of trimyristin, tripalmitin, tristearin and triolein	A.J. Bennett, M.A. Billett, A.M. Salter, E.H. Mangiapane, J.S. Bruce, K.L. Anderton, C.B. Marenah, N. Lawson and D.A. White	<b>167–173</b>
Triplex-DNA stabilization by hydralazine and the presence of anti-(triplex DNA) antibodies in patients treated with hydralazine	T.J. Thomas, J.R. Seibold, L.E. Adams and E.V. Hess	<b>183–188</b>
Selective loss of the uncoupling protein from light versus heavy mitochondria of brown adipocytes after a decrease in noradrenergic stimulation <i>in vivo</i> and <i>in vitro</i>	M.L. Bonet, F. Serra, J.C. Matamala, F.J. García-Palmer and A. Palou	<b>327–331</b>
<b>Membranes and bioenergetics</b>		
Major changes in Complex I activity in mitochondria from aged rats may not be detected by direct assay of NADH:coenzyme Q reductase	M.L. Genova, C. Castelluccio, R. Fato, G. Parenti Castelli, M. Merlo Pich, G. Formiggini, C. Bovina, M. Marchetti and G. Lenaz	<b>105–109</b>
<b>Receptors and signal transduction</b>		
Insulin and insulin-like growth factor I (IGF-I) stimulate GLUT4 glucose transporter translocation in <i>Xenopus</i> oocytes	S. Mora, P. Kaliman, J. Chillarón, X. Testar, M. Palacín and A. Zorzano	<b>59–65</b>
Quantification of signalling components and amplification in the $\beta$ -adrenergic-receptor-adenylate cyclase pathway in isolated adult rat ventricular myocytes	S.R. Post, R. Hilal-Dandan, K. Urasawa, L.L. Brunton and P.A. Insel	<b>75–80</b>
Naphthalenesulphonamides block neutrophil superoxide production by intact cells and in a cell-free system: is myosin light chain kinase responsible for these effects?	P.G. Heyworth, R.W. Erickson, J. Ding, J.T. Curnutte and J.A. Badwey	<b>81–87</b>
The protein phosphatase inhibitor calyculin A stimulates chemokine production by human synovial cells	N.J. Jordan, M.L. Watson and J. Westwick	<b>89–95</b>
Increase in receptor-like protein tyrosine phosphatase activity and expression level on density-dependent growth arrest of endothelial cells	F. Gaits, R.Y. Li, A. Ragab, J.M.F. Ragab-Thomas and H. Chap	<b>97–103</b>
Defective signal-transduction pathways in T-cells from autoimmune MRL- <i>lpr/lpr</i> mice are associated with increased polyamine concentrations	T.J. Thomas, U.B. Gunnia, J.R. Seibold and T. Thomas	<b>175–182</b>
Role of cytosolic phospholipase A <sub>2</sub> in arachidonic acid release of rat-liver macrophages: regulation by Ca <sup>2+</sup> and phosphorylation	P. Ambs, M. Baccarini, E. Fitzke and P. Dieter	<b>189–195</b>

---

Time-dependent effects of lithium on the agonist-stimulated accumulation of second messenger inositol 1,4,5-trisphosphate in SH-SY5Y human neuroblastoma cells	G.V. Los, I.P. Artemenko and L.E. Hokin	<b>225–232</b>
Structure and function of C-CAM1: effects of the cytoplasmic domain on cell aggregation	S.-H. Lin, W. Luo, K. Earley, P. Cheung and D.C. Hixson	<b>239–245</b>
Muscarinic stimulation exerts both stimulatory and inhibitory effects on the concentration of cytoplasmic Ca <sup>2+</sup> in the electrically excitable pancreatic B-cell	P. Gilon, M. Nenquin and J.-C. Henquin	<b>259–267</b>
Extracellular ATP activates different signalling pathways in rat Sertoli cells	C. Foresta, M. Rossato, P. Bordon and F. Di Virgilio	<b>269–274</b>
<b>Cell biology and development</b>		
Inhibition of asialoglycoprotein endocytosis and degradation in rat hepatocytes by protein phosphatase inhibitors	I. Holen, P.B. Gordon, P.E. Strømhaug, T.O. Berg, M. Fengsrud, A. Brech, N. Roos, T. Berg and P.O. Seglen	<b>317–326</b>
<b>BJ Letters</b>		
The DAG family of glycosyl hydrolases combines two previously identified protein families	B.H. Dagnall, I.T. Paulsen and M.H. Saier, Jr.	<b>349–350</b>
Families, superfamilies and subfamilies of glycosyl hydrolases	B. Henrissat and A. Romeu	<b>350–351</b>