

Research Communications

Bacterial expression and characterization of human secretory class V phospholipase A ₂	S.-K. Han, E.T. Yoon and W. Cho	353–357
Cloning of a functional promoter of the human sodium/iodide-symporter gene	M. Behr, T.L. Schmitt, C.R. Espinoza and U. Loos	359–363
Measurements of oxidative DNA damage by gas chromatography–mass spectrometry: ethanethiol prevents artifactual generation of oxidized DNA bases	A. Jenner, T.G. England, O.I. Aruoma and B. Halliwell	365–369
Glutathione transferase Zeta catalyses the oxygenation of the carcinogen dichloroacetic acid to glyoxylic acid	Z. Tong, P.G. Board and M.W. Anders	371–374

Research Papers

Proteins

Functional effects of single amino acid substitutions in the region of Phe ¹¹³ to Asp ¹³⁸ in the plasminogen activator inhibitor 1 molecule	G.-C. Sui and B. Wiman	409–415
Triplet structure of human von Willebrand factor	B.E. Fischer, K.B. Thomas, U. Schlokot and F. Dörner	483–488
Binding of a burst-phase intermediate formed in the folding of denatured D-glyceraldehyde-3-phosphate dehydrogenase by chaperonin 60 and 8-anilino-1-naphthalenesulphonic acid	X.-L. Li, X.-D. Lei, H. Cai, J. Li, S.-L. Yang, C.-C. Wang and C.-L. Tsou	505–511

Enzymes

Phosphinic peptides, the first potent inhibitors of astacin, behave as extremely slow-binding inhibitors	I. Yiallourous, S. Vassiliou, A. Yiotakis, R. Zwilling, W. Stöcker and V. Dive	375–379
pH-dependent inhibition by azide and fluoride of the iron superoxide dismutase from <i>Propionibacterium shermanii</i>	B. Meier, C. Scherk, M. Schmidt and F. Parak	403–407
The <i>dhnA</i> gene of <i>Escherichia coli</i> encodes a Class I fructose biphosphate aldolase	G.J. Thomson, G.J. Howlett, A.E. Ashcroft and A. Berry	437–445
Induction of matrix metalloproteinase activation cascades based on membrane-type 1 matrix metalloproteinase: associated activation of gelatinase A, gelatinase B and collagenase 3	S. Cowell, V. Knäuper, M.L. Stewart, M.-P. d'Ortho, H. Stanton, R.M. Hembry, C. López-Otín, J.J. Reynolds and G. Murphy	453–458
Purification and partial characterization of a peroxidase from plant cell cultures of <i>Cassia didymobotrya</i> and biotransformation studies	A. Vitali, B. Botta, G. Delle Monache, S. Zappitelli, P. Ricciardi, S. Melino, R. Petruzzelli and B. Giardina	513–519
Characterization of the leupeptin-inactivating enzyme from <i>Streptomyces exfoliatus</i> SMF13 which produces leupeptin	I.S. Kim, Y.B. Kim and K.J. Lee	539–545
Study of stereospecificity in mushroom tyrosinase	J.C. Espín, P.A. García-Ruiz, J. Tudela and F. García-Cánovas	547–551
Fixed-point methods for computing the equilibrium composition of complex biochemical mixtures	P. Kuzmič	571–575
Cystathionine γ -synthase from <i>Arabidopsis thaliana</i> : purification and biochemical characterization of the recombinant enzyme overexpressed in <i>Escherichia coli</i>	S. Ravanel, B. Gakière, D. Job and R. Douce	639–648
S-Nitrosoglutathione is a substrate for rat alcohol dehydrogenase class III isoenzyme	D.E. Jensen, G.K. Belka and G.C. Du Bois	659–668

Carbohydrates and lipids

Metabolism and apoptotic properties of elevated ceramide in HT29 ^{rev} cells	R.J. Veldman, K. Klappe, D. Hoekstra and J.W. Kok	563–569
---	---	----------------

Gene structure and expression

- Nucleolar protein p120 contains an arginine-rich domain that binds to ribosomal RNA 387–393
 W.C. Gustafson, C.W. Taylor, B.C. Valdez, D. Henning, A. Phippard, Y. Ren, H. Busch and E. Durban
- Regulation of type I collagen mRNA in lung fibroblasts by cystine availability 417–422
 D.C. Rishikof, P.-P. Kuang, C. Poliks and R.H. Goldstein
- Functional implications of ribosomal protein L2 in protein biosynthesis as shown by *in vivo* replacement studies 423–430
 M. Ühlein, W. Weglöhner, H. Urlaub and B. Wittmann-Liebold
- Cloning and characterization of a novel sequence-specific single-stranded-DNA-binding protein 447–452
 D. Bayarsaihan, R.J. Soto and L.N. Lukens
- Pyrazole-inducible proteins in DBA/2 mouse liver bind with high affinity to the 3'-untranslated regions of the mRNAs of coumarin hydroxylase (*CYP2A5*) and *c-jun* 473–481
 M. Thulke-Gross, M. Hergenahn, A. Tilloy-Ellul, M. Lang and H. Bartsch
- α - and δ -tocopherol induce expression of hepatic α -tocopherol-transfer-protein mRNA 577–581
 H. Fechner, M. Schlame, F. Guthmann, P.A. Stevens and B. Rüstow
- Structure of the human hexokinase type I gene and nucleotide sequence of the 5' flanking region 607–613
 A. Ruzzo, F. Andreoni and M. Magnani
- cDNA cloning reveals two mouse $\beta 5$ integrin transcripts distinct in cytoplasmic domains as a result of alternative splicing 631–637
 H. Zhang, S.M. Tan and J. Lu

Regulation of metabolism

- A distinct difference in the metabolic stimulus–response coupling pathways for regulating proinsulin biosynthesis and insulin secretion that lies at the level of a requirement for fatty acyl moieties 553–561
 R.H. Skelly, L.C. Bollheimer, B.L. Wicksteed, B.E. Corkey and C.J. Rhodes

Membranes and bioenergetics

- Subunit interaction of vacuolar H⁺-pyrophosphatase as determined by high hydrostatic pressure 395–402
 S.J. Yang, S.J. Ko, Y.R. Tsai, S.S. Jiang, S.Y. Kuo, S.H. Hung and R.L. Pan
- The reaction of halides with pulsed cytochrome *bo* from *Escherichia coli* 459–464
 A.J. Moody, C.S. Butler, N.J. Watmough, A.J. Thomson and P.R. Rich
- The negative charge of glutamic acid-820 in the gastric H⁺,K⁺-ATPase α -subunit is essential for K⁺ activation of the enzyme activity 465–472
 H.P.H. Hermsen, H.G.P. Swarts, J.B. Koenderink and J.J.H.H.M. De Pont
- Mevalonate-derived isopentenyl diphosphate is the biosynthetic precursor of ubiquinone prenyl side chain in tobacco BY-2 cells 615–621
 A. Disch, A. Hemmerlin, T.J. Bach and M. Rohmer

Receptors and signal transduction

- Expression of *v-src* in mammary epithelial cells induces transcription via STAT3 381–385
 P.D. Smith and M.R. Crompton
- cADP-ribose formation by blood platelets is not responsible for intracellular calcium mobilization 431–436
 P. Ohlmann, C. Leray, C. Ravanat, A. Hallia, D. Cassel, J.-P. Cazenave and C. Gachet
- Inhibition of kinases impairs neutrophil activation and killing of *Staphylococcus aureus* 489–495
 B. Schnyder, P.C. Meunier and B.D. Car
- Inactivating peptide of the *Shaker* B potassium channel: conformational preferences inferred from studies on simple model systems 497–504
 J.A. Encinar, A.M. Fernández, E. Gil-Martín, F. Gavilanes, J.P. Albar, J.A. Ferragut and J.M. González-Ros
- Activation of S6 kinase in human neutrophils by calcium pyrophosphate dihydrate crystals: protein kinase C-dependent and phosphatidylinositol-3-kinase-independent pathways 531–537
 C. Tudan, J.K. Jackson, L. Charlton, S.L. Pelech, B. Sahl and H.M. Burt
- Involvement of the Ras/extracellular signal-regulated kinase signalling pathway in the regulation of ERCC-1 mRNA levels by insulin 591–597
 W. Lee-Kwon, D. Park and M. Bernier
- Peroxynitrite modulates tyrosine phosphorylation and phosphoinositide signalling in human neuroblastoma SH-SY5Y cells: attenuated effects in human 1321N1 astrocytoma cells 599–606
 X. Li, P. De Sarno, L. Song, J.S. Beckman and R.S. Jope
- Histones and basic polypeptides activate Ca²⁺/cation influx in various cell types 623–630
 A. Gamberucci, R. Fulceri, P. Marcolongo, W.F. Pralong and A. Benedetti
- Purification of catalytic domain of rat spleen p72^{syk} kinase and its phosphorylation and activation by protein kinase C 649–657
 P. Borowski, M. Heiland, L. Kornetzky, S. Medem and R. Laufs

Cell biology and development

Species-specificity in endoplasmic reticulum signal peptide utilization revealed by proteins from <i>Trypanosoma brucei</i> and <i>Leishmania</i>	A. Al-Qahtani, M. Teilhet and K. Mensa-Wilmot	521–529
Two types of H ⁺ -ATPase are involved in the acidification of internal compartments in <i>Trypanosoma cruzi</i>	D.A. Scott and R. Docampo	583–589
Insulin targeting to the regulated secretory pathway after fusion with green fluorescent protein and firefly luciferase	A.E. Pouli, H.J. Kennedy, J.G. Schofield and G.A. Rutter	669–675

BJ Letters

Specificity of cysteine-rich domains in diacylglycerol kinases and protein kinases C	B. Houssa and W.J. van Blitterswijk	677–679
Hypothesis: ceramide conditionally activates atypical protein kinases C, Raf-1 and KSR through binding to their cysteine-rich domains	W.J. van Blitterswijk	679–680