

Review**Protein–protein interactions in intracellular
Ca²⁺-release channel function**
by J.J. Mackrill

345–361

Research Papers**Proteins**

A sensitive immunoassay for rat fatty acid translocase (CD36) using phage antibodies selected on cell transfectants: abundant presence of fatty acid translocase/CD36 in cardiac and red skeletal muscle and up-regulation in diabetes

M.M.A.L. Pelsers, J.T. Lutgerink, F.A. van Nieuwenhoven, N.N. Tandon, G.J. van der Vusse, J.-W. Arends, H.R. Hoogenboom and J.F.C. Glatz

407–414

Dermatopontin interacts with transforming growth factor β and enhances its biological activity

O. Okamoto, S. Fujiwara, M. Abe and Y. Sato

537–541

Isolation of a laminin-binding protein from the protozoan parasite *Leishmania donovani* that may mediate cell adhesion

A. Ghosh, K. Bandyopadhyay, L. Kole and P.K. Das

551–558

Enzymes

Structural aspects of the effectiveness of bisphosphonates as competitive inhibitors of the plant vacuolar proton-pumping pyrophosphatase

R. Gordon-Weeks, S. Parmar, T.G.E. Davies and R.A. Leigh

373–377

A novel 17 β -hydroxysteroid dehydrogenase in the fungus *Cochliobolus lunatus*: new insights into the evolution of steroid-hormone signalling

T. Lanišnik Rižner, G. Moeller, H.H. Thole, M. Žakej-Mavrič and J. Adamski

425–431

Novel peptide inhibitor of ecto-ADP-ribosyl cyclase of bone marrow stromal cell antigen-1 (BST-1/CD157)

A. Sato, S. Yamamoto, K. Ishihara, T. Hirano and H. Jingami

491–496

Substitution of Asp-309 by Asn in the Arg-Asp-Pro (RDP) motif of *Acetobacter diazotrophicus* levansucrase affects sucrose hydrolysis, but not enzyme specificity

F.R. Batista, L. Hernández, J.R. Fernández, J. Arrieta, C. Menéndez, R. Gómez, Y. Támara and T. Pons

503–506

Overexpression of the regulatory subunit of γ -glutamylcysteine synthetase in HeLa cells increases γ -glutamylcysteine synthetase activity and confers drug resistance

S.R. Tipnis, D.G. Blake, A.G. Shepherd and L.I. McLellan

559–566

Cloning and characterization of a simian UDP-glucuronosyltransferase enzyme UGT2B20, a novel C₁₉ steroid-conjugating protein

O. Barbier, A. Bélanger and D.W. Hum

567–574

L-ATP is recognized by some cellular and viral enzymes: does chance drive enzymic enantioselectivity?

A. Verri, A. Montecucco, G. Gosselin, V. Boudou, J.-L. Imbach, S. Spadari and F. Focher

585–590

Carbohydrates and lipids

Importance of splicing for prosaposin sorting

L. Madar-Shapiro, M. Pasmanik-Chor, A.M. Vaccaro, T. Dinur, A. Dagan, S. Gatt and M. Horowitz

433–443

Lipoarabinomannans: characterization of the multiacylated forms of the phosphatidyl-*myo*-inositol anchor by NMR spectroscopy

J. Nigou, M. Gilleron and G. Puzo

453–460

| | | |
|--|--|---------|
| Basic fibroblast growth factor does not prevent heparan sulphate proteoglycan catabolism in intact cells, but it alters the distribution of the glycosaminoglycan degradation products | S. Tumova, B.A. Hatch, D.J. Law and K.J. Bame | 471–481 |
| Gene structure and expression | | |
| Molecular characterization of the <i>Acremonium chrysogenum</i> <i>cefG</i> gene product: the native deacetylcephalosporin C acetyltransferase is not processed into subunits | J. Velasco, S. Gutierrez, S. Campoy and J.F. Martin | 379–385 |
| Interactions between the <i>Escherichia coli</i> cAMP receptor protein and the C-terminal domain of the α subunit of RNA polymerase at Class I promoters | E.C. Law, N.J. Savery and S.J.W. Busby | 415–423 |
| Transcription factor Sp1 activates involucrin promoter activity in non-epithelial cell types | E.B. Banks, J.F. Crish and R.L. Eckert | 507–512 |
| Occurrence and expression of members of the ferritin gene family in cowpeas | A.J. Wardrop, R.E. Wicks and B. Entsch | 523–530 |
| p21 ^{WAF1} expression by an activator of protein kinase C is regulated mainly at the post-transcriptional level in cells lacking p53: important role of RNA stabilization | M. Akashi, Y. Osawa, H.P. Koeffler and M. Hachiya | 607–616 |
| Regulation of metabolism | | |
| Formation of apolipoprotein-specific high-density lipoprotein particles from lipid-free apolipoproteins A-I and A-II | M.A. Clay, D.A. Cehic, D.H. Pyle, K.-A. Rye and P.J. Barter | 445–451 |
| <i>N</i> -Acetylsphingosine stimulates phosphatidylglycerolphosphate synthase activity in H9c2 cardiac cells | F.Y. Xu, S.L. Kelly and G.M. Hatch | 483–490 |
| Long-term maintenance of low concentrations of fructose for the study of hepatic glucose phosphorylation | J.W. Phillips, D.C. Henly and M.N. Berry | 497–501 |
| Glutamine synthesis is heterogeneous and differentially regulated along the rabbit renal proximal tubule | B. Ferrier, A. Conjard, M. Martin and G. Baverel | 543–550 |
| Membranes and bioenergetics | | |
| A comparative study of the activation of protein kinase C α by different diacylglycerol isomers | P. Sánchez-Piñera, V. Micol, S. Corbalán-García and J.C. Gómez-Fernández | 387–395 |
| Characterization of membrane polypeptides from pea leaf peroxisomes involved in superoxide radical generation | E. López-Huertas, F.J. Corpas, L.M. Sandalio and L.A. del Río | 531–536 |
| Receptors and signal transduction | | |
| Insulin and insulin-like growth factor I up-regulate <i>GLUT4</i> gene expression in fetal brown adipocytes, in a phosphoinositide 3-kinase-dependent manner | A.M. Valverde, P. Navarro, T. Teruel, R. Conejo, M. Benito and M. Lorenzo | 397–405 |
| Effects of transiently expressed atypical (ξ , λ), conventional (α , β) and novel (δ , ϵ) protein kinase C isoforms on insulin-stimulated translocation of epitope-tagged GLUT4 glucose transporters in rat adipocytes: specific interchangeable effects of protein kinases C- ξ and C- λ | G. Bandyopadhyay, M.L. Standaert, U. Kikkawa, Y. Ono, J. Moscat and R.V. Farese | 461–470 |
| Determination of the calcium-binding sites of the C2 domain of protein kinase C α that are critical for its translocation to the plasma membrane | S. Corbalán-García, J.A. Rodríguez-Alfaro and J.C. Gómez-Fernández | 513–521 |
| Role of phosphatidylinositol 3,4,5-trisphosphate in regulating the activity and localization of 3-phosphoinositide-dependent protein kinase-1 | R.A. Currie, K.S. Walker, A. Gray, M. Deak, A. Casamayor, C.P. Downes, P. Cohen, D.R. Alessi and J. Lucocq | 575–583 |
| Epidermal growth factor receptor activation is localized within low-buoyant density, non-caveolar membrane domains | M.G. Waugh, D. Lawson and J.J. Hsuan | 591–597 |

Epidermal growth factor increases the level of the cyclin-dependent kinase (CDK) inhibitor p21/CIP1 (CDK-interacting protein 1) in A431 cells by increasing the half-lives of the p21/CIP1 transcript and the p21/CIP1 protein

L.E. Johannessen, S.L. Knardal and I.H. Madhus

599–606

Cell biology and development

Differential expression of two novel Munc13 proteins in rat brain

I. Augustin, A. Betz, C. Herrmann, T. Jo and N. Brose

363–371