

**Review**

- The marks, mechanisms and memory of epigenetic states in mammals** 1–10  
*by V.K. Rakyan, J. Preis, H.D. Morgan and E. Whitelaw*

**Research Communication**

- Arg-Tyr-Asp (RYD) and Arg-Cys-Asp (RCD) motifs in dendroaspin promote selective inhibition of  $\beta_1$  and  $\beta_3$  integrins B. Wattam, D. Shang, S. Rahman, S. Egglezou, M. Scully, V. Kakkar and X. Lu 11–17

**Research Papers**

- Engineering of a glycosidase Family 7 cellobiohydrolase to more alkaline pH optimum: the pH behaviour of *Trichoderma reesei* Cel7A and its E223S/A224H/L225V/T226A/D262G mutant D. Becker, C. Braet, H. Brumer III, M. Claeysens, C. Divne, B.R. Fagerström, M. Harris, T.A. Jones, G.J. Kleywegt, A. Koivula, S. Mahdi, K. Piens, M.L. Sinnott, J. Ståhlberg, T.T. Teeri, M. Underwood and G. Wohlfahrt 19–30
- The potential mechanism in the diabetogenic action of streptozotocin: inhibition of pancreatic  $\beta$ -cell *O*-GlcNAc-selective *N*-acetyl- $\beta$ -D-glucosaminidase R.J. Konrad, I. Mikolaenko, J.F. Tolar, K. Liu and J.E. Kudlow 31–41
- Pterin interactions with distinct reductase activities of NO synthase M.M. Pantke, A. Reif, J.G. Valtschanoff, Z. Shutenko, A. Frey, R.J. Weinberg, W. Pfeleiderer and H.H.H.W. Schmidt 43–51
- Cloning of a lymphatic peptide/histidine transporter K. Sakata, T. Yamashita, M. Maeda, Y. Moriyama, S. Shimada and M. Tohyama 53–60
- Subcellular distribution of chelatable iron: a laser scanning microscopic study in isolated hepatocytes and liver endothelial cells F. Petrat, H. de Groot and U. Rauen 61–69
- Role of glycine-534 and glycine-1179 of human multidrug resistance protein (MDR1) in drug-mediated control of ATP hydrolysis G. Szakács, C. Özvegy, É. Bakos, B. Sarkadi and A. Váradí 71–75
- Induction of gene expression via activator protein-1 in the ascorbate protection against UV-induced damage M.V. Catani, A. Rossi, A. Costanzo, S. Sabatini, M. Levrero, G. Melino and L. Avigliano 77–85
- Heterodimerization of the epidermal-growth-factor (EGF) receptor and ErbB2 and the affinity of EGF binding are regulated by different mechanisms L.E. Johannessen, K.E. Haugen, A.C. Østfold, E. Stang and I.H. Madhus 87–96
- H<sub>c</sub> fragment (C-terminal portion of the heavy chain) of tetanus toxin activates protein kinase C isoforms and phosphoproteins involved in signal transduction C. Gil, I. Chaib-Oukadour, J. Blasi and J. Aguilera 97–103
- Inhibition of neuronal nitric oxide synthase by 6-nitrocatecholamines, putative reaction products of nitric oxide with catecholamines under oxidative stress conditions A. Palumbo, G. Astarita and M. d'Ischia 105–110
- Simple, sensitive and accurate method for the quantification of prothrombin mRNA by using competitive PCR P.K. Grover, A.M.F. Stapleton, K. Miyazawa and R.L. Ryall 111–120
- Human  $\alpha_1$ -acid glycoprotein binds to CCR5 expressed on the plasma membrane of human primary macrophages A. Atemezem, E. Mbemba, R. Vassy, H. Slimani, L. Saffar and L. Gattegno 121–128
- Human tear lipocalin acts as an oxidative-stress-induced scavenger of potentially harmful lipid peroxidation products in a cell culture system M. Lechner, P. Wojnar and B. Redl 129–135



Single-cell imaging of graded $\text{Ins}(1,4,5)P_3$ production following G-protein-coupled-receptor activation	M.S. Nash, K.W. Young, G.B. Willars, R.A.J. Challiss and S.R. Nahorski	<b>137–142</b>
Expression of a prenylation-deficient Rab4 inhibits the GLUT4 translocation induced by active phosphatidylinositol 3-kinase and protein kinase B	M. Cormont, N. Gautier, K. Ilc and Y. Le Marchand-Brustel	<b>143–149</b>
Molten-globule structure and membrane binding of the N-terminal protease-resistant domain (63–193) of the steroidogenic acute regulatory protein (StAR)	M. Song, H. Shao, A. Mujeeb, T.L. James and W.L. Miller	<b>151–158</b>
Crustacean hyperglycaemic hormone (CHH)-like peptides and CHH-precursor-related peptides from pericardial organ neurosecretory cells in the shore crab, <i>Carcinus maenas</i> , are putatively spliced and modified products of multiple genes	H. Dirksen, D. Böcking, U. Heyn, C. Mandel, J.S. Chung, G. Baggerman, P. Verhaert, S. Daufeldt, T. Plösch, P.P. Jaros, E. Waelkens, R. Keller and S.G. Webster	<b>159–170</b>
Transcriptional regulation of the <i>KEL</i> gene and Kell protein expression in erythroid and non-erythroid cells	V. Camara-Clayette, C. Rahuel, C. Lopez, C. Hattab, V. Verkarre, O. Bertrand and J.-P. Cartron	<b>171–180</b>
Barley arabinoxylan arabinofuranohydrolases: purification, characterization and determination of primary structures from cDNA clones	R.C. Lee, R.A. Burton, M. Hrmova and G.B. Fincher	<b>181–189</b>
Activation of store-mediated calcium entry by secretion-like coupling between the inositol 1,4,5-trisphosphate receptor type II and human transient receptor potential (hTrp1) channels in human platelets	J.A. Rosado and S.O. Sage	<b>191–198</b>
Isolation of ubiquitin–E2 (ubiquitin-conjugating enzyme) complexes from erythroleukaemia cells using immunoaffinity techniques	K. Takada, T. Hirakawa, H. Yokosawa, Y. Okawa, H. Taguchi and K. Ohkawa	<b>199–206</b>
Self-association and precursor protein binding of <i>Saccharomyces cerevisiae</i> Tom40p, the core component of the protein translocation channel of the mitochondrial outer membrane	D.M. Gordon, J. Wang, B. Amutha and D. Pain	<b>207–215</b>
Substitution of murine ferrochelatase glutamate-287 with glutamine or alanine leads to porphyrin substrate-bound variants	R. Franco, A.S. Pereira, P. Tavares, A. Mangravita, M.J. Barber, I. Moura and G.C. Ferreira	<b>217–222</b>
Inhibition of <i>Escherichia coli</i> CTP synthase by glutamate $\gamma$ -semialdehyde and the role of the allosteric effector GTP in glutamine hydrolysis	S.L. Bearne, O. Hekmat and J.E. MacDonnell	<b>223–232</b>
$\alpha 1$ Integrin cytoplasmic domain is involved in focal adhesion formation via association with intracellular proteins	K. Löster, D. Vossmeier, W. Hofmann, W. Reutter and K. Danker	<b>233–240</b>
Chimaeras reveal the role of the catalytic core in the activation of the plasma membrane $\text{Ca}^{2+}$ pump	W. Ba-Thein, A.J. Caride, A. Enyedi, K. Pászty, C.L. Croy, A.G. Filoteo and J.T. Penniston	<b>241–245</b>
Effects of the human papilloma virus HPV-16 E7 oncoprotein on glycolysis and glutaminolysis: role of pyruvate kinase type M2 and the glycolytic-enzyme complex	S. Mazurek, W. Zwerschke, P. Jansen-Dürr and E. Eigenbrodt	<b>247–256</b>
Cloning and identification of MYPT3: a prenylatable myosin targeting subunit of protein phosphatase 1	J.A. Skinner and A.R. Saltiel	<b>257–267</b>
Human oestrogenic $17\beta$ -hydroxysteroid dehydrogenase specificity: enzyme regulation through an NADPH-dependent substrate inhibition towards the highly specific oestrone reduction	A. Gangloff, A. Garneau, Y.-W. Huang, F. Yang and S.-X. Lin	<b>269–276</b>
Cation-dependent structural features of $\beta$ -casein-(1–25)	K.J. Cross, N.L. Huq, W. Bicknell and E.C. Reynolds	<b>277–286</b>
Phosphatidylinositol transfer proteins and protein kinase C make separate but non-interacting contributions to the phosphorylation state necessary for secretory competence in rat mast cells	J.A. Pinxteren, B.D. Gomperts, D. Rogers, S.E. Phillips, P.E.R. Tatham and G.M.H. Thomas	<b>287–296</b>