

Review**Mechanisms balancing skeletal matrix synthesis and degradation**

329–341

by H.C. Blair, M. Zaidi and P.H. Schlesinger

*Published on the Internet 24 May 2002, DOI 10.1042/BJ20020165***Research Communication**

Phosphorylation-dependent interaction of the synaptic vesicle proteins cysteine string protein and synaptotagmin I

G.J.O. Evans and A. Morgan

343–347

*Published as BJ Immediate Publication 4 April 2002, DOI 10.1042/BJ20020123***Research Papers**Metabolic control of resistance of human epithelial cells to H₂O₂ and NO stresses

C. Le Goffe, G. Vallette L. Charrier, T. Candelon, C. Bou-Hanna, J.-F. Bouhours and C.L. Laboissee

349–359

*Published on the Internet 24 May 2002, DOI 10.1042/BJ20011856*Peroxisome-proliferator-activated receptor- α (PPAR α) deficiency leads to dysregulation of hepatic lipid and carbohydrate metabolism by fatty acids and insulin

M.C. Sugden, K. Bulmer, G.F. Gibbons, B.L. Knight and M.J. Holness

361–368

*Published on the Internet 24 May 2002, DOI 10.1042/BJ20011699*Impaired noradrenaline-induced lipolysis in white fat of aP2-*Ucp1* transgenic mice is associated with changes in G-protein levels

P. Flachs, J. Novotný, F. Baumruk, K. Bardová, L. Bouřová, I. Mikšík, J. Šponarová, P. Svoboda and J. Kopecký

369–376

Published on the Internet 24 May 2002, DOI 10.1042/BJ20011438

Interaction of protein tyrosine phosphatase (PTP) 1B with its substrates is influenced by two distinct binding domains

S. Dadke and J. Chernoff

377–383

*Published on the Internet 24 May 2002, DOI 10.1042/BJ20011372*Control mechanisms operating for lipid biosynthesis differ in oil-palm (*Elaeis guineensis* Jacq.) and olive (*Olea europaea* L.) callus cultures

U.S. Ramli, D.S. Baker, P.A. Quant and J.L. Harwood

385–391

Published on the Internet 24 May 2002, DOI 10.1042/BJ20010202

Control analysis of lipid biosynthesis in tissue cultures from oil crops shows that flux control is shared between fatty acid synthesis and lipid assembly

U.S. Ramli, D.S. Baker, P.A. Quant and J.L. Harwood

393–401

Published on the Internet 24 May 2002, DOI 10.1042/BJ20010203

Fibrinogen is degraded and internalized during incubation with neutrophils, and fibrinogen products localize to electron lucent vesicles

R. Kirsch, M.A. Jaffer, V.E. Woodburne, T. Sewell, S.L. Kelly, R.E. Kirsch and E.G. Shephard

403–412

Published on the Internet 24 May 2002, DOI 10.1042/BJ20011406

Gene promoter of apoptosis inhibitory protein IAP2: identification of enhancer elements and activation by severe hypoxia

Z. Dong, J. Nishiyama, X. Yi, M.A. Venkatachalam, M. Denton, S. Gu, S. Li and M. Qiang

413–421

*Published on the Internet 24 May 2002, DOI 10.1042/BJ20011431*Mycolic acid biosynthesis and enzymic characterization of the β -ketoacyl-ACP synthase A-condensing enzyme from *Mycobacterium tuberculosis*

L. Kremer, L.G. Dover, S. Carrère, K.M. Nampoothiri, S. Lesjean, A.K. Brown, P.J. Brennan, D.E. Minnikin, C. Locht and G.S. Besra

423–430

Published on the Internet 24 May 2002, DOI 10.1042/BJ20011628

Membrane depolarization-induced contraction of rat caudal arterial smooth muscle involves Rho-associated kinase

M. Mita, H. Yanagihara, S. Hishinuma, M. Saito and M.P. Walsh

431–440

Published on the Internet 24 May 2002, DOI 10.1042/BJ20020191

Ascorbate up-regulates MLH1 (Mut L homologue-1) and p73: implications for the cellular response to DNA damage <i>Published on the Internet 24 May 2002, DOI 10.1042/BJ20011713</i>	M.V. Catani, A. Costanzo, I. Savini, M. Levrero, V. De Laurenzi, J.Y.J. Wang, G. Melino and L. Avigliano	441–447
Retinoic acid receptor-related orphan receptor (ROR) α 4 is the predominant isoform of the nuclear receptor ROR α in the liver and is up-regulated by hypoxia in HepG2 human hepatoma cells <i>Published on the Internet 24 May 2002, DOI 10.1042/BJ20011558</i>	C. Chauvet, B. Bois-Joyeux and J.-L. Danan	449–456
Structure–function analysis of yeast piD261/Bud32, an atypical protein kinase essential for normal cell life <i>Published on the Internet 24 May 2002, DOI 10.1042/BJ20011376</i>	S. Facchin, R. Lopreiato, S. Stocchetto, G. Arrigoni, L. Cesaro, O. Marin, G. Carignani and L.A. Pinna	457–463
Natural variation in the extent of phosphorylation of bone phosphoproteins as a function of <i>in vivo</i> new bone formation induced by demineralized bone matrix in soft tissue and bony environments <i>Published on the Internet 24 May 2002, DOI 10.1042/BJ20011272</i>	E. Salih, J. Wang, J. Mah and R. Fluckiger	465–474
Polarized cholesterol and phospholipid efflux in cultured gall-bladder epithelial cells: evidence for an ABCA1-mediated pathway <i>Published on the Internet 24 May 2002, DOI 10.1042/BJ20011493</i>	J. Lee, A. Shirk, J.F. Oram, S.P. Lee and R. Kuver	475–484
Studies on the relevance of the glycan at Asn-52 of the α -subunit of human chorionic gonadotropin in the $\alpha\beta$ dimer <i>Published on the Internet 24 May 2002, DOI 10.1042/BJ20011482</i>	P.J.A. Erbel, S.R. Haseley, J.P. Kamerling and F.G. Vliegenthart	485–495
Biochemical characterization and subcellular localization of human copper transporter 1 (hCTR1) <i>Published on the Internet 24 May 2002, DOI 10.1042/BJ20011803</i>	A.E.M. Klomp, B.B.J. Tops, I.E.T. van den Berg, R. Berger and L.W.J. Klomp	497–505
The terminal six amino-acids of the carboxy cytoplasmic tail of CD36 contain a functional domain implicated in the binding and capture of oxidized low-density lipoprotein <i>Published on the Internet 24 May 2002, DOI 10.1042/BJ20011373</i>	E. Malaud, D. Hourton, L.M. Giroux, E. Ninio, R. Buckland and J.L. McGregor	507–515
IgA Fc receptor (Fc α R) cross-linking recruits tyrosine kinases, phosphoinositide kinases and serine/threonine kinases to glycolipid rafts <i>Published on the Internet 24 May 2002, DOI 10.1042/BJ20011696</i>	M.L. Lang, Y.-W. Chen, L. Shen, H. Gao, G.A. Lang, T.K. Wade and W.F. Wade	517–525
Purification of the <i>Escherichia coli</i> ammonium transporter AmtB reveals a trimeric stoichiometry <i>Published on the Internet 24 May 2002, DOI 10.1042/BJ20011761</i>	D. Blakey, A. Leech, G.H. Thomas, G. Coutts, K. Findlay and M. Merrick	527–535
Oxygen-evoked changes in transcriptional activity of the 5'-flanking region of the human amiloride-sensitive sodium channel (α ENaC) gene: role of nuclear factor κ B <i>Published on the Internet 24 May 2002, DOI 10.1042/BJ20011651</i>	D.L. Baines, M. Janes, D.J. Newman and O.G. Best	537–545
Increased actin polymerization reduces the inhibition of serum response factor activity by Yin Yang 1 <i>Published on the Internet 24 May 2002, DOI 10.1042/BJ20020269</i>	P.D. Ellis, K.M. Martin, C. Rickman, J.C. Metcalfe and P.R. Kemp	547–554
Sterol 14 α -demethylase activity in <i>Streptomyces coelicolor</i> A3(2) is associated with an unusual member of the CYP51 gene family <i>Published on the Internet 24 May 2002, DOI 10.1042/BJ20011380</i>	D.C. Lamb, K. Fowler, T. Kieser, N. Manning, L.M. Podust, M.R. Waterman, D.E. Kelly and S.L. Kelly	555–562
Activation of mouse Pi-class glutathione S-transferase gene by Nrf2 (NF-E2-related factor 2) and androgen <i>Published on the Internet 24 May 2002, DOI 10.1042/BJ20011756</i>	H. Ikeda, M.S. Serria, I. Kakizaki, I. Hatayama, K. Satoh, S. Tsuchida, M. Muramatsu, S. Nishi and M. Sakai	563–570
Repression of Smad2 and Smad3 transactivating activity by association with a novel splice variant of CCAAT-binding factor C subunit <i>Published on the Internet 24 May 2002, DOI 10.1042/BJ20011703</i>	F. Chen, K. Ogawa, X. Liu, T.M. Stringfield and Y. Chen	571–577

Reactive sulphur species: an <i>in vitro</i> investigation of the oxidation properties of disulphide S-oxides <i>Published on the Internet 24 May 2002, DOI 10.1042/BJ20011882</i>	G.I. Giles, K.M. Tasker, C. Collins, N.M. Giles, E. O'Rourke and C. Jacob	579–585
Mutational analysis of the carbohydrate-binding activity of the NeuAc(α -2,6)Gal/GalNAc-specific type 2 ribosome-inactivating protein from elderberry (<i>Sambucus nigra</i>) fruits <i>Published on the Internet 24 May 2002, DOI 10.1042/BJ20020006</i>	Y. Chen, P. Rouge, W.J. Peumans and E.J.M. van Damme	587–592